

Research Article**Assessment of anti-obesity activity of Tamarindus leaves on butter induced hyperlipidemia in mice****Geetha Rani Valaparla*, Gutta Vishnu Sai, Devireddysudheer Kumar***Department of Pharmacology, A.M. Reddy memorial College of Pharmacy, Petlurivaripalem, Narasaraopet, 522601 Guntur, Andhra Pradesh, India*

Received: 27 April 2022

Revised: 4 May 2022

Accepted: 13 June 2022

Abstract

Objective: Obesity is one of the world's leading problems, which is major cause of Cardiovascular Diseases. During COVID - 19 pandemic many young children and all age group people are getting challenged by obesity due stagnant life style. To treat such life threatening disorder natural remedies acquire prior position, Plant produce a good deal of secondary metabolites which have a benefited mankind in various ways, including treatment of diseases The present research work was focused on developing a natural remedy using *Tamarindus indica*, an ancient plant with number of proven activities in Traditional medicine. **Materials and Methods:** In the literature survey, it was found that flavonoids, sterols, tannins and alkaloids shown promising effects to tackle obesity by various mechanisms, *Tamarindus indica* has shown the presence of saponins, flavonoids, glycosides, amino acids, alkaloids. With the above evidence, this plant has been selected for screening of its antiobesity activity against high fat diet induced obesity in mice. Studies were conducted using aqueous extract of *Tamarindus indica* leaves (AETI) on high fat diet (butter) induced model of hyperlipidemia in mice. **Results and Discussion:** During 21 days time period AETI low and high doses were induced to respective animal groups along with butter where Atorvastatin has been taken as standard drug. Butter enhanced the Cholesterol and triglyceride, LDL levels. At the end of 21 days blood samples were collected from the animal through retroperitoneal route. Biochemical parameters of AETI have been revealed its antiobesity activity.

Keywords: Obesity, AETI, high fat diet, antiobesity, Atorvastatin

Introduction

According to WHO overweight and obesity are defined as abnormal or excessive fat accumulation that presents a risk to health. A body mass index (BMI) over 25 is considered overweight, and over 30 is obese (Hassan et al., 2017).

Overweight and obesity are major risk factors for a number of chronic diseases, including cardiovascular diseases such as stroke, diabetes and its associated conditions including blindness, limb amputations, and the need for dialysis, musculoskeletal disorders including osteoarthritis. Obesity is also associated with some cancers, including endometrial, breast, ovarian, prostate, liver, gallbladder, kidney and colon. The risk of these non-communicable diseases increases even when a person is only slightly overweight and grows more

serious as the body mass index (BMI) climbs. WHO defines over weight as a BMI equal to or more than 25, obesity as a BMI equal to or more than 30.

Statins are a class of prescription medicines that have been used for decades to lower low density lipoprotein (LDL-C) cholesterol in the blood (Crebonell and Freire, 2005). Medicines in the statin class include atorvastatin, flustatin, lovastatin, pitavastatin, pravastatin, rosuvastatin and simvastatin.

Atorvastatin is a high efficacy and one of the most widely used lipid lowering medication (Jing et al., 2012; Goa et al., 2013). The efficacy and safety of Atorvastatin have been testified by more than 200 randomized controlled trails, with most sufficient clinical evidence among all statins (Adams et al., 2012). Atorvastatin is a reversible and competitive inhibitor of 3- hydroxy-3- methylglutaryl – coenzyme A reductase, decreasing the denovo cholesterol synthesis (Cong et al, 2015), has been selected as standard drug for the present research.

Plant produce a good deal of secondary metabolites which have a benefited mankind in various ways, including

***Address for Corresponding Author:**

Geetha Rani Valaparla
Associate Professor
Department of Pharmacology,
A.M. Reddy memorial College of Pharmacy, Petlurivaripalem,
Narasaraopet, 522601 Guntur, Andhra Pradesh, India
E mail: purnageetha15@gmail.com

DOI: <https://doi.org/10.31024/ajpp.2022.8.3.5>2455-2674/Copyright © 2022, N.S. Memorial Scientific Research and Education Society. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

treatment of diseases (Elain et al., 2002), *Tamarindus indica* belongs to the family Fabaceae and sub family Detarioideae. The plant has been used for centuries as a medicinal plant, fruit part of the plant has been more valuable, have often been reported in several pharmacopoeias. Leaves of the plant have good levels of proteins, fat, fiber and some vitamins such as thiamine, riboflavin, nicin, ascorbic acid, B- carotene (El-siddig et al., 2006) The leaves have proven hepatoprotective activity (Joeux et al., 1995), antimicrobial activity (Muthu et al., 2005), antifungal (Zohrameena et al., 2017; Khare, 2004; Melendez and Carriles, 2006; Lansc, 2007), antiseptic (Zohrameena et al, 2017; Khare,2004; Melendez and Carriles, 2006; Lansc, 2007), wound healing activity (Fabiya et al., 1993), anti emetic activity (Rasheed et al., 2015), antihistaminic activity etc (Jayade et al., 2010).

In the literature survey, it was found that flavanoids, sterols, tannins and alkaloids shown promising effects to tackle obesity by various mechanisms (Rohit, 2013), *Tamarindus indica* has shown the presence of saponins, flavanoids, glycosides, amino acids, alkaloids. With the above evidence, this plant has been selected for screening of its antiobesity activity against high fat diet induced obesity in mice.

Materials and Methods

Plant material collection and authentication

The plant material of *Tamarindus indica* leaves used for the investigation was collected from A M Reddy Memorial College of Pharmacy premises, Narasaraopet, Guntur Dt. The plant was identified and authenticated by P.SATYANARAYANA RAJU Department of botany from Acharya Nagarjuna University, Nagarjuna Nagar, Guntur, 522510, Guntur Dt, Andhra Pradesh.

Preparation of extract

The leaves of *Tamarindus indica* were collected and shade dried at room temperature and grinded coarsely before extraction. The leaves were extracted by hot maceration by using distilled water. The resulting extract was collected into air tight container. Thus, the prepared extract was used for further pharmacological evaluation.

Experimental Animals

Healthy male adult albino mice stains 20- 30 gms were selected for the study, obtained from Hyderabad. The animals were housed properly under 12:12 hours light and dark cycle and fed with proper food and water. National CPCSEA guidelines were strictly followed and all the studies were approved by the Institutional animal ethical committee.

Preliminary phytochemical evaluation

The aqueous extract of *Tamarindus indica* leaves was subjected to preliminary phytochemical analysis to assess the presence of

various phytoconstituents. It revealed the presence of saponins, flavanoids, glycosides, amino acids, alkaloids. All these tests were performed at A.M. Reddy Memorial College of pharmacy, Narasaraopet, Guntur, Andhra Pradesh.

Obesity Inducer

For the present study Butter was mainly used to induce obesity in animals. 400 mg of butter /kg body weight dissolved in 10ml of buffered saline was administered.

Test drug preparation

The two doses of test (AETI low dose and high dose) and Standard drug Atorvastatin are soluble in water, so were dissolved in distilled water and administered to the respective animal groups per oral. All the test concentrations were prepared freshly before administering to the animals. Butter was administered after 30min of test drug administration to all groups.

Animal groups

Group-1 was considered as control group which received only the vehicle.

Group-2 was considered as the high fat diet group which received the butter.

Group-3 was considered as first testing group and received the test extract that is aqueous extract of *Tamarindus indica* (AETI) 250 mg per kg weight per oral along with butter.

Group-4 was considered as second test group and received the test extracts that is aqueous extracts of *Tamarindus indica* (AETI) 500mg per kg weight per oral along with butter.

Group-5 was considered as standard group which received the standard drug atorvastatin (10mg/ kg) along with butter.

Body weight determination Body weight of the animals was recorded every week before administering of the test drugs.

Sample collection: At the end of 21st day blood was collected from the retro orbital plexus after overnight fasting and then was centrifuged and the serum was obtained and was estimated for the total cholesterol, LDL, VLDL and HDL.

Biochemical Parameters

On the 21st day of the methodology that is after the last test drug administration, the mice were anesthetized with Diethyl ether and the blood samples were collected from retro orbital puncture using capillary tube into clotting tubes. Serum samples were analysed at Jayanti laboratory at Jayanti Hospital, Palnadu Road, Vinukonda, Guntur, Andhra Pradesh.

Statistical analysis

All data are expressed as mean \pm S.E.M. The data of all the experiments were analyzed using one way analysis of variance (ANOVA) followed by Tukey's post hoc test. In all the tests, criterion for statistical significance was $P < 0.05$

Results and Discussion

Obesity is the major and health problem in India and developing countries, which lead to important risk factors like atherosclerosis, stroke etc. Obesity increases the likelihood of other diseases. Obesity evokes the damages in various tissues, which in turn, deregulate the cellular functions leading to damage to various pathological conditions.

Atorvastatin is a high efficacy and one of the most widely used lipid lowering medication. The efficacy and safety of Atorvastatin have been testified by more than 200 randomized controlled trails, with most sufficient clinical evidence among all statins (Adams et al., 2012), Atorvastatin is a reversible and competitive inhibitor of 3-hydroxy-3-methylglutaryl- coenzyme A reductase, decreasing the denovo cholesterol synthesis (Cong et al., 2015), regulate LDL, VLDL-C levels has been selected as standard drug for the present research.

In the literature survey, it was found that flavanoids, sterols, tannins and alkaloids shown promising effects to tackle obesity by various mechanisms (Jerry, 2011), *Tamarindus indica* has shown the presence of saponins, flavanoids, glycosides, amino acids, alkaloids. With the above evidence, this plant has been selected for screening of its antiobesity activity against high fat diet induced obesity in mice. All the results of are given in table 1 and 2.

Table 1. Preliminary photochemical tests

S.NO	Photochemical tests	Inference
1	Test for alkaloids	+ve
2	Test for flavonoids	+ve
3	Test for glycosides	+ve
4	Test for amino acids	+ve
5	Test for saponins	+ve

Table 2. Effect of *Tamarindus indica* on bio-chemical parameters

Groups	Total cholesterol (mg/dl)	Triglycerides (mg/dl)	HDL (mg/dl)	LDL (mg/dl)	VLDL (mg/dl)
Group I (Control)	156 \pm 1.56	112.0 \pm 49.58	62.0 \pm 1.58	90 \pm 44	25 \pm 1.45
Group II (Butter Induced)	244.50 \pm 1.48	360.0 \pm 1.60	41.0 \pm 1.62	170 \pm 1.75	78.16 \pm 1.68
Group III (AETI 0.5ml)	200.90 \pm 1.58	190 \pm 1.60	44 \pm 1.58	110 \pm 1.68	37 \pm 1.60
Group IV (AETI 1 MI)	181 \pm 1.26	166.0 \pm 1.48	58.0 \pm 1.58	87 \pm 1.67	33 \pm 1.25
Group V (Std Atorvastatin)	161.50 \pm 1.94	162.54 \pm 1.58	38.48 \pm 1.41	120 \pm 0.60	41 \pm 1.53

High cholesterol is also referred to as hypercholesterolemia. Cholesterol is a fatty substance that is important part of the outer lining of cells in the body of the animals. Cholesterol is also found in the blood circulation of humans. It is also a precursor for the synthesis of steroid hormones. High levels of cholesterol can increase the risk of heart disease. Cholesterol is synthesized in all animal tissue. Increased amount of cholesterol leads to cardiovascular diseases particularly coronary heart disease (CHC) (Nordestgaard and Varbo, 2014). Total Cholesterol levels of AETI high dose were recorded as 181 \pm 1.26 and standard drug shown 161.50 \pm 1.26, in general controlling of cholesterol by AETI 161 \pm 1.26 is said as good reduction.

Triglycerides are a type of fat in the blood stream and in fat tissue. These are small enough to enter into the arterial wall and thus have the potential to accumulate and cause atherosclerosis (Nordestgaard and Varbo, 2014; Nordestgaard et al., 1995; Goldberg et al., 2011), Looking at TG level of both AETI and standard drug are almost similar 166.0 \pm 1.48 and 162.54 \pm 1.58, here just 4 points difference was observed.

HDL-C are inversely correlated with clinical events resulting from atherosclerosis, instead HDL protects against cardiovascular disease by regulating cholesterol efflux from tissue and modulating inflammation (Mohamad et al., 2011). In case of HDL-C, AETI 58.0 \pm 1.58 could not reach the value of Standard drug 38.48 \pm 1.41. So the study was ensuring to lower the HDL-C of AETI in a protecting way.

LDL is a risk factor and places a role at several steps of atherosclerosis. LDL is called low density lipoprotein because LDL particles tend to the less dense than other kinds of cholesterol particles. Increase in evidence as revealed that the concentration and the size of the LDL particles more powerfully relates to the degree of atherosclerosis progression than the concentration of cholesterol contained within all the LDL particles. Perhaps, it was observed that the test drug AETI 87 \pm 1.67 is able to bring down the LDL level very much than standard drug 120 \pm 0.60.

VLDL (very low density lipoprotein) transports cholesterol and triglycerides within the body, and are associated with

increased risk of coronary heart disease. It was observed good reduction in VLDL level with AETI (33 ± 1.25) than standard drug 41 ± 1.53 .

Conclusion

In congruence with these results, it may be confirmed that due to the presence of phytoconstituents such as saponins, flavanoids, glycosides, amino acids, alkaloids in the aqueous extract, it could be responsible for the observed significant anti-obesity activity. So, the present study proven that the aqueous extract of *Tamarindus indica* leaves exhibited significant Anti-obesity activity.

Acknowledgement

All the authors express their gratitude to A.M. Reddy memorial College of Pharmacy for valuable support.

Conflict of interest: None

References

- Adams SP, Tsang M, James MW. 2012. Lipid lowering efficacy of atorvastatin. Cochrane Database Systematic Reviewer. 12.
- Aparna Berteri R. 2003. Risk of coronary artery heart disease. Health Screen 1:28-29.
- Balentine R.J. 2011. Facts of obesity. Medicinenet 1:1-12.
- Cong ye Y, Liang X, Yang S. 2015. Use of Atorvastatin in lipid disorders and cardiovascular disease in Chinese Patients. Chinese medical journal. 128(2):259–266.
- Crebonell T, Freire E. 2005. Binding thermodynamics of ststins to hmg-coa reductage. Biochemistry. 44:11741-8.
- Elaine MS, Ana BR, Olindo A, Gazinelli G. 2002. Screening and fractionation of plant extracts with antiproliferative activity on human peripheral blood mononuclear Cells. Memorias do Instituto Oswaldo Cruz 97(8):1207-1212.
- El-Siddig G, Gunasena H, Prasad BA, Pushpakumara DKNP, Ramana KVR. 2006. *Tamarindus indica*. South ampton UK centre for under utilised crops.
- Fabiyi JP, Kela SL, Tal MK, Istifanus WA. 1993. Traditional therapy of dracunculiasis in the state of Bauchi Nigeria. Dakar Medical (38):193-195.
- Goa F, Zouh Y, Hu D, Ying X. 2013. Contemporary management and attainment of cholesterol targets for patients with dyslipidemia in China. PLoS One 8(4):7681.
- Goldberg IJ, Robert HE, Ruth Mc P. 2011. Triglycerides and heart disease: still a hypothesis? Arterioscler Thromb. Vascular Biology. 15(4):534-542.
- Gundamaraju R. 2013. Evaluation of Anti-Obesity Activity of *Lantana camara* Var Linn. by Progesterone Induced Obesity on Albino Mice. International Journal of Pharmacognosy and Phytochemical Research, 4(4):213-218.
- Hassan DA, Abdullah DAI-S, Tariq AA, Bassam AA, Kaled AA. 2017. Relationship between Body mass index and Obesity awareness in school students. Journal of Clinical Medicin Research, 9(6).
- Joyeux M, Francois M, Jacques F. 1995. Screening of antiradical, anti-lipoperoxidant and hepatoprotective effects of nine plant extracts used in Garibbean folk medicine. Phytotherapy Research, 9(3):228-230.
- Khare CP. 2004. Editor Encyclopaedia of Indian medicinal plant-Rational Western therapy.. Ayurvedic and other traditional usage. Botany, Springer Verlay.
- Lans C. 2007. Comparison of plants used for skin and stomach problems in Trinidad and Tobago with Asian ethnomedicine. Journal of Ethnobiology and Ethnomedicine, (3):102-109.
- Li J, PingChen Y, Li X, Armitage J, Feng F. 2012. Use of secondary preventive medications in patients with atherosclerotic disease in urban China. A cross-sectional study of 16, 860 patients. Chinese Medical Journal, 25:4361–4367.
- Meléndez PA, Carriles VA. 2006. Antibacterial properties of tropical plants from Puerto Rico. Phytomedicine, 13:272-276.
- Mohamad N, Reddy S, Brian JVL, Alem MF. 2011. HDL and cardiovascular disease atherogenic and athero protective mechanisms: Macmillan Publishers Limited. 8(4):222-32.
- Muthu SE, nandakumar S, Rao UA. 2005. The effect of methanolic extract of *Tamarindus indica* on the growth of clinical isolates of *Burkholderia pseudomallei*. Indian Journal of Medical Research, (8):122:525.
- Nordestgaard BG, Rechar W, Lewis B. 1995. Selective retention of VLDL, IDL, and LDL in the arterial intima of genetically hyperlipidemic rabbits in vivo. Molecular size as a determinant of fractional loss from the intima-inner media. Arterioscler Thromb Vasc Biology, 15:534-42.
- Nordestgaard BG, Varbo A. 2014. Triglycerides and cardiovascular disease. Lancet, 384:626-635.
- Rasheed AH, Amna SK, Hasan SK, Khalid, Hassan Al. 2015. Acute and subacute toxicity of of *Tamarindus indica* L fruit methanolic extract. Worls Journal of Pharmaceutical Research, 4(12):1519-1527.
- Tayade P, Bordi SN, Jagtab SA, VP Patel, G Vaishnav, YR Girbane, Swapnil C, Geeshan SK. 2010. Effect of *Tamarindus indica* Linn. Against isolated goat tracheal and guinea pig ileum preparation. International Journal of Comprehensive Pharmacy, 2(6).
- Zohrameena S, Paramepeed B, Maad K, Mohd M. 2017. Medicinal uses & pharmacological activity of *Tamarindus indica*. World Journal of Pharmaceutical Science, 5(2):121-133.